



World Health Organization


Mortality Country Fact Sheet 2006

Philippines



Summary	Year	Males	Females	Both sexes
Population (millions)	2005	42	41	83
Life expectancy (years)	2004	65	72	68
Under-5 mortality (per 1 000 live births)	2004	40	28	34
Adult mortality (per 1 000)	2004	269	149	
Maternal mortality (per 100 000 live births)	2000		200	

Source: World Health Statistics 2006




Legend:


- WHO Western Pacific Region
- Philippines

Source: World Health Statistics 2006

Under-5 mortality



Source: i) WHO mortality database ii) World Health Statistics 2006




Note: rate for 5 years preceding the survey
Source: DHS 1993, DHS 1998, DHS 2003

Under-5 mortality: for highest and lowest quintiles
Philippines, DHS 2003

Wealth/assets quintiles	Lowest	Highest	Ratio
	66	21	3.1
Sex	Males	Females	Ratio
	48	34	1.4
Urban/Rural	Rural	Urban	Ratio
	52	30	1.7
Mother's education quintiles	None	Higher	Ratio
	105	29	3.7

Note: rate per 1 000 live births for 10-year period preceding the survey
Source: DHS 2003



Note: rate for 10-year period preceding the survey
Source: DHS 2003

Causes of death in children under-5

Distribution of causes of death among children under 5 years of age Philippines, 2000-2003			Annual estimated proportions of death by cause for neonates Philippines, 2000		
Causes	Deaths ^b		Causes	Deaths ^c	
	(%)	Regional average (%)		(%)	Regional average ^c (%)
Total neonatal deaths	100	100	Total neonatal deaths	100	100
Neonatal causes ^a	37	47	Neonatal tetanus	1	3
HIV/AIDS	0	0	Severe infection ^a	17	21
Diarrhoeal diseases	12	12	Birth asphyxia	28	26
Measles	1	1	Diarrhoeal diseases	1	1
Malaria	0	0	Congenital anomalies	8	8
Pneumonia	13	14	Preterm birth ^b	37	32
Injuries	3	7	Others	8	8
Others	33	18			

a. Includes diarrhoea during neonatal period
b. Sum of individual proportions may not add up to 100% due to rounding.

a. Includes deaths from pneumonia, meningitis, sepsis/septicaemia and other infections during the neonatal period.
b. Includes only deaths directly attributed to prematurity and to specific complications of preterm birth such as surfactant deficiency, but not all deaths in preterm infants.
c. Sum of individual proportions may not equal 100% due to rounding.

Causes of Death

Top ten causes of death, all ages Philippines, 2002				Life expectancy at birth among males (years) Philippines, 2004	
Causes	Deaths		Years of Life Lost	30 40 50 60 70 80 90 100	
	(000)	(%)		[Scale]	
All causes	448	100	100	[Scale]	
Lower respiratory infections	46	10	11	[Scale]	
Ischaemic heart disease	45	10	6	[Scale]	
Tuberculosis	44	10	8	[Scale]	
Hypertensive heart disease	31	7	4	[Scale]	
Perinatal conditions	25	6	11	[Scale]	
Cerebrovascular disease	24	5	3	[Scale]	
Violence	16	4	5	[Scale]	
Diarrhoeal diseases	12	3	5	[Scale]	
Diabetes mellitus	11	3	2	[Scale]	
Chronic obstructive pulmonary disease	9	2	1	[Scale]	

Source: [Death and DALY estimates by cause, 2002](http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls)
http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls

Legend:
 WHO Western Pacific Region
 Philippines

Source: World Health Statistics 2006